arXiv:1605.03422v1 [math.GM] 6 May 2016

Curves with normal planes at constant distance from a fixed point

Yasemin ALAGOZ

Department of Mathematics, Faculty of Arts and Sciences, Yildiz Technical University Istanbul-TURKEY

Email: ygulluk@yildiz.edu.tr, yasemingulluk@yahoo.com

May 12, 2016

Abstract

We consider the curves whose all normal planes are at the same distance from a fixed point and obtain some characterizations of them in the 3-dimensional Euclidean space.

Keywords: Curve, plane curve, distance, spherical curve, involute.

2000 Mathematics Subject Classification: 53A04.

1 Introduction

Beng-Yen Chen investigated the curves whose position vectors lie in their rectifying planes in the 3-dimensional Euclidean space, [1]. These curves are called the rectifying curves which are not plane curves. They can be equivalently defined as the twisted curves whose osculating planes are at the same distance from a fixed point. Here we consider the curves whose all normal planes are at the same distance from a fixed point and obtain some characterizations of them in the 3-dimensional Euclidean space.

We first review the results on the rectifying curves in the 3-dimensional Euclidean space, [1].

Let C be a unit-speed curve with the position vector $\mathbf{r}(s)$. Therefore s is the natural parameter of C. Frenet formulas follow as

$$\mathbf{t}' = \kappa\mathbf{n}, \quad \mathbf{n}' = -\kappa\mathbf{t} + \tau\mathbf{b}, \quad \mathbf{b}' = -\tau\mathbf{n} \tag{1}$$

where $\mathbf{t}, \mathbf{n}$ and $\mathbf{b}$ are Frenet vectors and κ the curvature and τ the torsion of

the curve.
A rectifying curve can be defined by the equation

$$< \mathbf{r}, \mathbf{n} > = 0 \tag{2}$$

By means of (1) we can write it as

$$< \mathbf{r}, \mathbf{t}' > = 0 \tag{3}$$

Accordingly $< \mathbf{r}, \mathbf{t} >' = < \mathbf{r}, \mathbf{t}' > + < \mathbf{t}, \mathbf{t} > = 1$ and so

$$< \mathbf{r}, \mathbf{t} > = \int_0^s 1 ds = s \tag{4}$$

Integrating (4) we have

$$< \mathbf{r}, \mathbf{r} > = \rho^2 = s^2 + c \tag{5}$$

where c is a constant. Without loss the generality we can write (5) as

$$\rho^2 = s^2 + a^2 \tag{6}$$

where a is a positive number.
(2) , (4) and (6) imply that the position vector can be written as

$$\mathbf{r} = s\mathbf{t} + a\mathbf{b} \tag{7}$$

Differentiating (7) we have

$$\mathbf{t} = \mathbf{t} + s(\kappa \mathbf{n}) + a(-\tau \mathbf{n})$$

and so

$$s\kappa = a\tau \tag{8}$$

This implies that a rectifying curve is a twisted curve, that is $\tau \neq 0$.
On the other hand, from (7) we have

$$< \mathbf{r}, \mathbf{b} > = a \tag{9}$$

So if a curve is a rectifying curve, then its osculating planes are at the same distance from the origin. Conversely, let us assume that osculating planes of a twisted curve are at the same distance from the origin. Differentiating (9) we get (2) and so it is a rectifying curve. Therefore we have the following property:

A twisted curve is a rectifying curve if and only if its osculating planes are at the same distance from a fixed point.

On the other hand, if a unit spherical curve C_1 is defined by the equation

$$\mathbf{r}_1(s) = \frac{\mathbf{r}(s)}{\sqrt{s^2 + a^2}} \tag{10}$$

where $\mathbf{r}(s)$ is the position vector of a rectifying curve C with the natural parameter s, it can be show that the position vector of a rectifying curve C can be represented as

$$\mathbf{r}(s_1) = (a \sec s_1)\mathbf{r}_1(s_1) \tag{11}$$

where s_1 is the natural parameter of the spherical curve C_1. Then the tangent unit vector of the curve C can be written as

$$\mathbf{t}(s_1) = (\sin s_1)\mathbf{r}_1(s_1) + (\cos s_1)\mathbf{t}_1(s_1) \tag{12}$$

where $\mathbf{t}_1(s_1)$ is the tangent unit vector of the spherical curve C_1. Differentiating the last equation we have

$$\kappa \mathbf{n} a \sec^2 s_1 = (\cos s_1)(\mathbf{r}_1 + \kappa_1 \mathbf{n}_1) \tag{13}$$

κ_1 is the curvature of the spherical curve C_1.
From (13) we find the following relation between the curvatures of the curves C and C_1 associated with each other:

$$a^2\kappa^2 = (\cos s_1)^6(\kappa_1^2 - 1) \tag{14}$$

The last relation can be written, in terms of the natural parameter s of the rectifying curve, as

$$(s^2 + a^2)^3\kappa^2 = a^4(\kappa_1^2 - 1) \tag{15}$$

2 Curves with normal planes at the same distance from a fixed point

Let us assume that all normal planes of a curve C with the position vector $\mathbf{r} = \mathbf{r}(s)$ are at the same distance from a fixed point where s is the natural parameter of C. We denote Frenet vectors and the curvature and the torsion of C by $\mathbf{t}, \mathbf{n}$ and $\mathbf{b}$ and κ and τ respectively. We can choose the fixed point as the origin. Therefore our condition becomes

$$< \mathbf{r}, \mathbf{t} > = c_1 = \text{constant} \tag{16}$$

By the integration we find that

$$< \mathbf{r}, \mathbf{r} >= \rho_1^2 = 2c_1 s + c_2 \tag{17}$$

where c_2 is a constant.
Since the case of $c_1 = 0$ corresponds to a spherical curve, we can assume that $c_1 \neq 0$. Then, without loss the generality, we can write (17) as

$$< \mathbf{r}, \mathbf{r} >= \rho_1^2 = 4\epsilon c^2 s, \qquad \epsilon = \text{sign(s)} \tag{18}$$

where c is a positive constant. Then (16) reduces to

$$< \mathbf{r}, \mathbf{t} > = 2\epsilon c^2 \tag{19}$$

(18) and (19) imply

$$\epsilon s \geq c^2 \tag{20}$$

Differentiating (19) we have

$$\kappa < \mathbf{r}, \mathbf{n} >= -1 \tag{21}$$

or

$$< \mathbf{r}, \mathbf{n} > = -R \tag{22}$$

where $R = 1/\kappa$ is the radius of curvature.
From (19) and (22) we can write the position vector as

$$\mathbf{r} = 2\epsilon c^2 \mathbf{t} - R\mathbf{n} + A\mathbf{b} \tag{23}$$

Differentiating (23) we have

$$\mathbf{t} = \mathbf{t} + (2\epsilon c^2 \kappa - R' - A\tau)\mathbf{n} + (A' - R\tau)\mathbf{b} \tag{24}$$

and

$$A\tau = 2\epsilon c^2 \kappa - R', \qquad A' = R\tau \tag{25}$$

Therefore we first have

$$RR' = 2\epsilon c^2 \Longleftrightarrow \tau = 0 \tag{26}$$

So the curve is a plane curve if and only if

$$RR' = 2\epsilon c^2 \tag{27}$$

Without the generality, for case of a plane curve we can write

$$\mathbf{r} = 2\epsilon c^2 \mathbf{t} + (-R)\mathbf{n} \tag{28}$$

Then from (18) and (28)

$$R^2 = 4c^2(\epsilon s - c^2) \tag{29}$$

Accordingly the natural equations of the curve are

$$\kappa = \frac{1}{2c\sqrt{\epsilon s - c^2}}, \qquad \tau = 0 \tag{30}$$

We can now assume that our curve is a twisted curve, that is $\tau \neq 0$. Then

$$A = 2\epsilon c^2 \kappa T - R'T, \qquad A' = R\tau \tag{31}$$

where $T = \frac{1}{\tau}$ is the radius of torsion. Hence (23) can be written as

$$\mathbf{r} = (2\epsilon c^2)\mathbf{t} + (-R)\mathbf{n} + (2\epsilon c^2 \kappa T - R'T)\mathbf{b} \tag{32}$$

Therefore, according to (18) we have

$$4\epsilon c^2 s = 4c^4 + R^2 + (2\epsilon c^2 \kappa T - R'T)^2 \tag{33}$$

On the other hand, from (31) we can write

$$R\tau + (R'T)' - 2\epsilon c^2 (\kappa T)' = 0 \tag{34}$$

We can show that (33) and (34) are equivalent equations. In fact, the equation (34) can be written as

$$[R\tau + (R'T)' - 2\epsilon c^2 (\kappa T)'][RR' - 2\epsilon c^2] = 0 \tag{35}$$

because of $RR' - 2\epsilon c^2 = 0$, corresponds to the plane curve. Then we have

$$2\epsilon c^2 = [R\tau + (R'T)' - 2\epsilon c^2 (\kappa T)']R'T + 4c^4 \kappa T(\kappa T)' - 2\epsilon c^2 \kappa T(R'T)' \tag{36}$$

and

$$4c^2 = 2RR' + 2(2\epsilon c^2 \kappa T - R'T)[2\epsilon c^2 (\kappa T)' - (R'T)'] \tag{37}$$

This implies that

$$4\epsilon c^2 = (R^2)' + [(2\epsilon c^2 \kappa T - R'T)^2]' \tag{38}$$

Therefore we have

$$R^2 + (2\epsilon c^2 \kappa T - R'T)^2 = \int_{c^2}^{\epsilon s} 4c^2 dt = 4\epsilon c^2 s - 4c^4$$

It is obvious that using the equation (33) we obtain the equation (34).

3 Spherical curves associated with a curve with normal planes at constant distance from a point

Let us define a unit spherical curve C_1 by the equation

$$\mathbf{r}_1(s) = \frac{\mathbf{r}(s)}{2c\sqrt{\epsilon s}} \tag{39}$$

where $\mathbf{r}(s)$ is the position vector of a curve with normal planes at constant distance from a fixed point.
So we have

$$\mathbf{r}(s) = 2c\sqrt{\epsilon s}\mathbf{r}_1(s) \tag{40}$$

Differentiating we get

$$\mathbf{t} = \frac{\epsilon c}{\sqrt{\epsilon s}}\mathbf{r}_1(s) + (2c\sqrt{\epsilon s})\mathbf{r}_1'(s) \tag{41}$$

From the last equation we find

$$|\mathbf{r}_1'(s)| = \frac{\sqrt{\epsilon s - c^2}}{2c\epsilon s} = \frac{ds_1}{ds} \tag{42}$$

for the speed of the spherical curve. And so the natural parameter of the unit spherical curve C_1, from $s_1 = \epsilon \int_{c^2}^{\epsilon s} \frac{\sqrt{t-c^2}}{2ct} dt$ is obtained as

$$s_1 = \epsilon\left(\frac{\sqrt{\epsilon s - c^2}}{c} - \arctan\frac{\sqrt{\epsilon s - c^2}}{c}\right) \tag{43}$$

Since

$$\frac{ds_1}{ds} = \frac{\sqrt{\epsilon s - c^2}}{2c\epsilon s} > 0 \tag{44}$$

there exists a function $s(s_1)$ which satisfies the equation (43) Therefore (39) can be written as

$$\mathbf{r}_1(s(s_1)) = \frac{\mathbf{r}(s(s_1))}{2c\sqrt{\epsilon s(s_1)}} \tag{45}$$

So for a given curve C with normal planes at constant distance from a fixed point, whose position vector is $\mathbf{r}(s_1) = \mathbf{r}(s(s_1))$, we have a unit spherical curve C_1 whose position vector $\mathbf{r}_1(s_1) = \mathbf{r}_1(s(s_1))$ is defined by (45) with the natural parameter s_1. We call C_1 the unit spherical curve associated with the curve C with normal planes at constant distance from a fixed point.

Now let us consider a curve C defined by

$$\mathbf{r}(s_1) = (2c\sqrt{\epsilon s(s_1)}\mathbf{r}_1(s_1) \tag{46}$$

where $\mathbf{r}_1(s_1)$ is the position vector of a unit spherical curve C_1 with the natural parameter s_1 and $s(s_1)$ is defined by (43) and c is constant. Let $\mathbf{t}_1,\ \mathbf{n}_1,\ \mathbf{b}_1$ and κ_1 and τ_1 be Frenet vectors and the curvature and the torsion of C_1 respectively. According to (43)

$$s' = \frac{ds}{ds_1} = \frac{2c\epsilon s}{\sqrt{\epsilon s - c^2}} \tag{47}$$

Differentiating (46) with respect to s_1 we have

$$\mathbf{r}' = \frac{d\mathbf{r}}{ds_1} = 2\epsilon c^2 \frac{\sqrt{\epsilon s}}{\sqrt{\epsilon s - c^2}}\mathbf{r}_1 + (2c\sqrt{\epsilon s})\mathbf{t}_1 \tag{48}$$

Since $< \mathbf{r}_1, \mathbf{r}_1 >= 1,\ < \mathbf{r}_1, \mathbf{t}_1 >= 0$ and $< \mathbf{t}_1, \mathbf{t}_1 >= 1$ from (48) we have

$$|\mathbf{r}'| = \frac{2c\epsilon s}{\sqrt{\epsilon s - c^2}}$$

So the unit tangent vector of C is found as

$$\mathbf{t}(s_1) = \frac{\epsilon c}{\sqrt{\epsilon s}}\mathbf{r}_1(s_1) + \frac{\sqrt{\epsilon s - c^2}}{\sqrt{\epsilon s}}\mathbf{t}_1(s_1) \tag{49}$$

Since $< \mathbf{r}, \mathbf{r}_1 >= 2c\sqrt{\epsilon s}$ and $< \mathbf{r}, \mathbf{t}_1 >= 0$

$$< \mathbf{r}, \mathbf{t} > = 2\epsilon c^2 \tag{50}$$

This means that the curve C is a curve with normal planes at constant distance from a fixed point. We call C the curve with normal planes at constant distance from a fixed point associated with the unit spherical curve C_1.
Let us now differentiate (49) with respect to s. We have

$$\kappa\mathbf{n} = -\frac{c}{2\epsilon s\sqrt{\epsilon s}}\mathbf{r}_1 + \frac{1}{2\sqrt{\epsilon s}\sqrt{\epsilon s - c^2}}\mathbf{t}_1 + \frac{\epsilon s - c^2}{2\epsilon c s\sqrt{\epsilon s}}\kappa_1\mathbf{n}_1 \tag{51}$$

Using $< \mathbf{n}, \mathbf{n} > = 1, < \mathbf{r}_1, \mathbf{r}_1 >= 1, < \mathbf{t}_1, \mathbf{t}_1 >= 1, < \mathbf{n}_1, \mathbf{n}_1 >= 1, < \mathbf{r}_1, \mathbf{t}_1 >= 0, < \mathbf{t}_1, \mathbf{n}_1 >= 0$ and $\kappa_1 < \mathbf{r}_1, \mathbf{n}_1 >= -1$, from (51) we obtain the following relation between curvatures of the curves C and C_1 associated with each other:

$$\kappa^2 = \frac{3s^2 - 3c^2\epsilon s + c^4}{4\epsilon s^3(s - c^2)} + \frac{(\epsilon s - c^2)^2}{4c^2\epsilon s^3}\kappa_1^2 \tag{52}$$

or

$$\kappa^2 = \frac{1}{4c^2(\epsilon s - c^2)} + \frac{(\epsilon s - c^2)^2}{4c^2\epsilon s^3}(\kappa_1^2 - 1) \tag{53}$$

Since $\kappa_1 < \mathbf{r}_1, \mathbf{n}_1 >= -1$,the curvature of the unit spherical curve is not smaller than 1, that is $\kappa_1 \geq 1$.

Example 1

The equation (53) implies that the curvature of the unit spherical curve associated with the plane curve given by the natural equations (30) is $\kappa_1 = 1$. This means that the spherical curve is a great circle of the unit sphere. Hence we can obtain the cartesian equations of the plane curve associated with a great circle of the unit sphere using the equation (40). In fact, we can choose the equation of a great circle of the unit sphere as

$$\mathbf{r}_1 = (\cos s_1, \sin s_1, 0) \tag{54}$$

Since $s_1 = \epsilon(\frac{\sqrt{\epsilon s - c^2}}{c} - \arctan\frac{\sqrt{\epsilon s - c^2}}{c})$,

$$\cos s_1 = \cos\frac{\sqrt{\epsilon s - c^2}}{c}\frac{c}{\sqrt{\epsilon s}} + \epsilon\sin\frac{\sqrt{\epsilon s - c^2}}{c}\frac{\sqrt{\epsilon s - c^2}}{\sqrt{\epsilon s}}$$

and

$$\sin s_1 = \epsilon\sin\frac{\sqrt{\epsilon s - c^2}}{c}\frac{c}{\sqrt{\epsilon s}} - \cos\frac{\sqrt{\epsilon s - c^2}}{c}\frac{\sqrt{\epsilon s - c^2}}{\sqrt{\epsilon s}}$$

Then from (40) we obtain the Cartesian equations of the plane curve with normal

planes at constant distance from the origin as

$$\begin{aligned} x &= 2c^2 \cos\frac{\sqrt{\epsilon s - c^2}}{c} + 2\epsilon c\sqrt{\epsilon s - c^2}\sin\frac{\sqrt{\epsilon s - c^2}}{c} \\ y &= 2\epsilon c^2 \sin\frac{\sqrt{\epsilon s - c^2}}{c} - 2c\sqrt{\epsilon s - c^2}\cos\frac{\sqrt{\epsilon s - c^2}}{c} \\ z &= 0 \end{aligned} \tag{55}$$

This curve is a plane curve with normal planes at a distance $2c^2$ from the origin. This means that all normal lines of the curve are at a distance $2c^2$ from the origin. So it is an involute of the circle of radius $2c^2$ centered at the origin. Since it is an involute of a plane curve, it is also a plane curve [3, p,88]. In fact, the position vector of an involute C of the curve C_2 whose position vector is $\mathbf{r}_2(s_2)$can be written as

$$\mathbf{r} = \mathbf{r}_2(s_2) + (c_2 - s_2)\mathbf{t}_2(s_2) \tag{56}$$

where c_2 is constant and $\mathbf{t}_2(s_2)$ is unit tangent vector to the curve C_2, [3, p.69], [5, p.40], [4, p.99]. Then

$$\mathbf{r}'{=}(c_2 - s_2)\kappa_2\mathbf{n}_2 \tag{57}$$

where κ_2 is the curvature of C_2 and $\mathbf{n}_2$ the principal normal vector. Therefore the unit tangent vector t to C can be written as

$$\mathbf{t} = -\epsilon_2\mathbf{n}_2, \qquad -\epsilon_2 = \text{sign}(c_2 - s_2) \tag{58}$$

Since C_2 is a circle of radius $2c^2$ centered at the origin,

$$\mathbf{r}_2 = (2c^2\cos\frac{s_2}{2c^2}, 2c^2\sin\frac{s_2}{2c^2}, 0) \tag{59}$$

$$\mathbf{t}_2 = (-\sin\frac{s_2}{2c^2}, \cos\frac{s_2}{2c^2}, 0) \tag{60}$$

$$\mathbf{n}_2 = (-\cos\frac{s_2}{2c^2}, -\sin\frac{s_2}{2c^2}, 0) \tag{61}$$

So $< \mathbf{r}_2, \mathbf{t}_2 >= 0$, $< \mathbf{r}_2, \mathbf{n}_2 >= -2c^2$. Then from (56) and (58) we have

$$< \mathbf{r}, \mathbf{t} >= 2\epsilon_2 c^2 \tag{62}$$

This means that the involute C of the circle C_2 is a plane curve with normal planes at a distance $2c^2$ from the origin.

Figure 1

Figure 1 illustrates the plane curve (55) with normal planes at a distance $2c^2 = 2(1)^2 = 2$ from the origin (the solid line for $\epsilon = 1$, the dashed line for $\epsilon = -1$). It is an involute of the circle of radius $2c^2 = 2$ centered at the origin. Let us note the equations (55) can be also obtained using the equations (56)-(61) or using (55) from the equations of a plane curve given by

$$\mathbf{r} = \mathbf{r}(s) = (x(s), y(s), 0) = (\int \cos(\int \kappa(s)ds)ds, \int \sin(\int \kappa(s)ds)ds, 0) \quad (63)$$

where s is the natural parameter of the curve, [3, p.87], [2, p.28], [4, p.99].
In the following we give an example of a twisted curve with normal planes at constant distance from a fixed point.

Example 2

Let us choose the unit spherical curve C_1 as the circle of radius $\frac{1}{\sqrt{2}}$ given by the equation

$$\mathbf{r}_1 = (\frac{1}{\sqrt{2}}\cos(\sqrt{2}s_1), \frac{1}{\sqrt{2}}\sin(\sqrt{2}s_1), \frac{1}{\sqrt{2}}) \quad (64)$$

Then the curve C associated with C_1 is given by the equation

$$\mathbf{r}(s) = (c\sqrt{2}\sqrt{\epsilon s}\cos(\sqrt{2}s_1), c\sqrt{2}\sqrt{\epsilon s}\sin(\sqrt{2}s_1), c\sqrt{2}\sqrt{\epsilon s}) \quad (65)$$

where $s_1 = \epsilon(\frac{\sqrt{\epsilon s - c^2}}{c} - \arctan\frac{\sqrt{\epsilon s - c^2}}{c})$.
Since the curvature of the circle C_1 is $\kappa_1 = \sqrt{2}$, (53) implies that

$$\kappa^2 = \frac{1}{4c^2(\epsilon s - c^2)} + \frac{(\epsilon s - c^2)^2}{4c^2\epsilon s^3} = \frac{\epsilon s^3 + (\epsilon s - c^2)^3}{4c^2\epsilon s^3(\epsilon s - c^2)} \quad (66)$$

Figure 2 illustrates the twisted curve (65) with normal planes at a distance $2c^2 = 2(1)^2 = 2$ from the origin (the solid line for $\epsilon = 1$, the dashed line for $\epsilon = -1$).

Figure 2

4 Curves of constant curvature with normal planes at the same distance from a fixed point

If $\kappa = \frac{1}{r}$ =constant, (32) reduce to

$$\mathbf{r} = (2\epsilon c^2)\mathbf{t} + (-r)\mathbf{n} + 2\frac{\epsilon c^2}{r}T\mathbf{b} \tag{67}$$

From (67) we have $< \mathbf{r}, \mathbf{n} > = -r$ =constant. This means that rectifying planes of a curve of constant curvature with normal planes at constant distance from a fixed point are also at constant distance from the same point.

Conversely if normal planes and rectifying planes of a curve are at constant distance from a fixed point, it is a curve of constant curvature. In fact by above conditions we can write

$$< \mathbf{r}, \mathbf{t} > = c_1 = \text{constant} \tag{68}$$

and

$$< \mathbf{r}, \mathbf{n} > = c_2 = \text{constant} \tag{69}$$

Differentiating (68) we have

$$1 + < \mathbf{r}, \kappa\mathbf{n} > = 0$$

and so

$$< \mathbf{r}, \mathbf{n} > = -R \tag{70}$$

Because of (69) and (70) the curve is a curve of constant curvature.

On the other hand for a curve of constant curvature with normal planes at constant distance from a fixed point (33) reduces to

$$4\epsilon c^2 s = 4c^4 + r^2 + \frac{4c^4}{r^2} T^2$$

Therefore the natural equations of such a curve are

$$R = r = \text{constant}, \qquad T^2 = r^2 \frac{2\epsilon h s - r^2 - h^2}{h^2}$$

where $h = 2c^2$ is the distance of the normal planes from the origin and r is the distance of the rectifying planes from the origin.

References

[1] Chen BY. When does the position vector of a space curve always lie in its rectifying plane?. Am Math Mon 110 2003; VOL: 147-152

[2] Eisenhart LP. A Treatise on The Differential Geometry of Curves and Surfaces. New York: Dover Publications, Inc, 1909.

[3] Goetz A. Introduction to Differential Geometry. California, London, Ontario: Addison Wesley Publication Company, Reading, Massachusetts, 1970.

[4] Hsiung CC. A First Course in Differential Geometry. Cambridge, MA 02238-2872 USA: International Press, P.O. Box 2872 , 1997

[5] Struik DJ. Lectures on Classical Differential Geometry. Cambridge,42, Mass: Addison-Wesley Press, Inc, 1950.

This figure "untitled2.png" is available in "png" format from:

http://arxiv.org/ps/1605.03422v1

This figure "untitled4.png" is available in "png" format from:

http://arxiv.org/ps/1605.03422v1